arXiv:1204.4892v1 [math.GT] 22 Apr 2012

On the Iwasawa invariants of a link in the 3-sphere

Teruhisa Kadokami Yasushi Mizusawa

Abstract. Based on the analogy between knots and primes, J. Hillman, D. Matei and M. Morishita defined the Iwasawa invariants for sequences of cyclic covers of links with an analogue of Iwasawa's class number formula of number fields. In this paper, we consider the existence of covers of links with prescribed Iwasawa invariants, discussing analogies in number theory. We also propose and consider a problem analogous to Greenberg's conjecture.

1. Introduction

Let $L = K_1 \cup K_2 \cup \cdots \cup K_r$ be an r-component link in the 3-sphere S^3. Let $\ell_{ij} = \mathrm{lk}(K_i, K_j)$ be the linking number of the components K_i and K_j $(i \neq j)$. Let X be the complement of an open tubular neighborhood of L, and $G_L = \pi_1(X)$ the link group of L. Let $\pi : \widetilde{X} \to X$ be the maximal abelian cover of X. Let $[m_i] \in G_L$ be the class of the meridian m_i of the component K_i, where we regard m_i as a loop from the base point $*$ of X. Then $H_1(X;\mathbb{Z}) \simeq \mathrm{Aut}(\widetilde{X}/X) \simeq G_L/G_L'$ is generated by $t_i = [m_i]G_L'$ $(i = 1, 2, \cdots, r)$ as a free $\mathbb{Z}$-module of rank r, where G_L' is the commutator subgroup of G_L. Put the Laurent polynomial ring $\Lambda_r = \mathbb{Z}[t_1^{\pm 1}, t_2^{\pm 1}, \cdots, t_r^{\pm 1}] \simeq \mathbb{Z}[G_L/G_L']$. Then the Alexander module $A_L = H_1(\widetilde{X}, \pi^{-1}(*);\mathbb{Z})$ and the link module $B_L = H_1(\widetilde{X};\mathbb{Z}) \simeq G_L'/G_L''$ are finitely generated Λ_r-module, where G_L'' is the commutator subgroup of G_L'.

Based on the analogies between link groups and Galois groups of number fields with restricted ramification (cf. [18] [19] [20] etc.), we regard $\widetilde{X}$ as an analogue of the maximal free abelian pro-p-extension $\widetilde{k}$ of a number field k, where p is a fixed prime number. Then the Iwasawa theory for $\widetilde{k}/k$ is set up in parallel with Alexander-Fox theory (cf. e.g. [4] [21]). Let $S = \{\wp_1, \wp_2, \cdots, \wp_r\}$ be the set of all primes of k lying over p, and $G_S = G_S(k)$ the Galois group of the maximal pro-p-extension of k unramified outside S. Then $H_1(G_S;\mathbb{Z}_p) \simeq G_S/G_S' \simeq \mathrm{Gal}(\widetilde{k}/k) \simeq \mathbb{Z}_p^{r_2+1+\delta}$ for a subgroup G_S' and $\delta \geq 0$, where $\mathbb{Z}_p$ is (the additive group of) the p-adic integer ring, and r_2 is the number of infinite complex places of k. Leopoldt's conjecture predicts that $\delta = 0$ for any k and p (cf. [27] Theorem 13.4). For the simplicity, we assume that $\delta = 0$ throughout this paper. If k has the class number $h_k \not\equiv 0 \pmod{p}$, G_S/G_S' is generated by $\tau_i G_S'$ $(i = 1, 2, \cdots, r_2+1)$ where each τ_i is an element of inertia group

---

2000 Mathematics Subject Classification: 57M25, 11R23.
Key words: Iwasawa invariant, Alexander polynomial, link module.

of some $\wp_j$. The S-ramified Iwasawa module $\mathfrak{X}_S = G'_S/G''_S$ is a finitely generated $\mathbb{Z}_p[[G_S/G'_S]]$-module, where G''_S is the commutator subgroup of G'_S.

$$\begin{array}{ccc}
\text{link } L = K_1 \cup K_2 \cup \cdots \cup K_r & \longleftrightarrow & \text{primes } S = \{\wp_1, \wp_2, \cdots, \wp_r\} \\
\text{link group } G_L & \longleftrightarrow & \text{Galois group } G_S \\
G_L/G'_L \simeq \mathrm{Aut}(\widetilde{X}/X) \simeq \mathbb{Z}^r & \longleftrightarrow & G_S/G'_S \simeq \mathrm{Gal}(\widetilde{k}/k) \simeq \mathbb{Z}_p^{r_2+1} \\
\text{meridians } [m_i] & \longleftrightarrow & \text{inertia groups } \langle \cdots, \tau_i, \cdots \rangle \\
\text{link module } B_L \simeq G'_L/G''_L & \longleftrightarrow & \text{Iwasawa module } \mathfrak{X}_S = G'_S/G''_S
\end{array}$$

For a surjective homomorphism $\sigma_{\mathbf{z}} : H_1(X;\mathbb{Z}) \to \mathbb{Z}$ (resp. $G_S/G'_S \to \mathbb{Z}_p$), we have an infinite cyclic cover $X_{\mathbf{z}} \to X$ (resp. a $\mathbb{Z}_p$-extension k_∞/k) corresponding to the kernel. Iwasawa [9] gave a formula for the growth of the p-parts of the class numbers in the $\mathbb{Z}_p$-extension k_∞/k (cf. §2). Hillman, Matei and Morishita [7] (and [10]) gave an analogous formula for the growth of the p-parts of the order of 1-homology groups in the tower of p^n-fold cyclic branched covering spaces of L along $X_{\mathbf{z}}$ (cf. §2 Theorem 2.1 (2)). These formulas are described by the Iwasawa invariants λ, μ and ν, which are closely related to the structure of link modules or Iwasawa modules.

In this paper, we show some basic properties of Iwasawa invariants of L, discussing analogies with Iwasawa theory of number fields. In §2 and §3, we decide the parity of λ-invariants of L and the existence of L with prescribed λ- and μ-invariants. We also give some examples of 2-component links L with explicit Iwasawa invariants. In §4 and §5, we propose and consider an analogous problem to Greenberg's conjecture which asserts that the maximal unramified quotient of $\mathfrak{X}_S$ is *pseudonull*.

Notations. For a noetherian unique factorization domain Λ and a finitely generated Λ-module A, we denote by $\mathrm{rank}_\Lambda A$ the free rank of A. The *divisorial hull* of an ideal E is the intersection of all principal ideals of Λ containing E (cf. [6] p.49). Let $E^{(i)}_\Lambda(A)$ be the ith elementary ideal of A, and $\Delta^{(i)}_\Lambda(A)$ a generator of the divisorial hull of $E^{(i)}_\Lambda(A)$ as a principal ideal. A finitely generated torsion Λ-module A is *pseudonull* if and only if the divisorial hull of the annihilator ideal $\mathrm{Ann}_\Lambda A$ is Λ (cf. [6] Theorem 3.5), i.e., $\mathrm{Ann}_\Lambda A$ has at least two relatively prime elements. We denote by $|A|$ the order of a $\mathbb{Z}$-module A. Note that $|A| = 0$ if A is infinite. Let v_p be the additive p-adic valuation normalized as $v_p(p) = 1$, and $\Phi_{p^n}(t) = \sum_{i=0}^{p-1}(t^{p^{n-1}})^i = (t^{p^n} - 1)/(t^{p^{n-1}} - 1)$ the p^nth cyclotomic polynomial for $n \geq 1$.

2. Iwasawa invariants and Alexander polynomials

For $f(t) \in \Lambda_1 = \mathbb{Z}[t^{\pm 1}]$, we denote by $\deg' f(t)$ the reduced degree as a Laurent polynomial. The Alexander polynomial of L is defined as

$$\Delta_L = \Delta_L(t_1, t_2, \cdots, t_r) = \Delta^{(1)}_{\Lambda_r}(A_L) = \Delta^{(0)}_{\Lambda_r}(B_L)$$

up to multiplication by units in Λ_r (cf. [12] Proposition 7.3.4 (2)). For $\mathbf{z} = (z_1, z_2, \cdots, z_r) \in \mathbb{Z}^r$ such that $\gcd(z_1, z_2, \cdots, z_r) = 1$ and $\prod_{i=1}^{r} z_i \neq 0$, we define a surjective homomorphism

$$\sigma_{\mathbf{z}} : H_1(X; \mathbb{Z}) \to \mathbb{Z} : t_i \mapsto z_i.$$

Then $\operatorname{Ker} \sigma_{\mathbf{z}}$ corresponds to an infinite cyclic cover $X_{\mathbf{z}} \to X$, and the reduced Alexander polynomial associated to $X_{\mathbf{z}}$ is defined in $\Lambda_1 \simeq \mathbb{Z}[\operatorname{Aut}(X_{\mathbf{z}}/X)]$ as

$$\Delta_{L,\mathbf{z}}(t) = \Delta^{(0)}_{\Lambda_1}(H_1(X_{\mathbf{z}}; \mathbb{Z})) = (t-1)\Delta_L(t^{z_1}, t^{z_2}, \cdots, t^{z_r})$$

if $r \geq 2$, and $\Delta_{L,\mathbf{z}}(t) = \Delta_L(t)$ if $r = 1$ (cf. [12] Proposition 7.3.10 (1)). For $\mathbf{1} = (1, 1, \cdots, 1)$, $X_{\mathbf{1}}$ is called the total linking number covering space of X. Let $M_{\mathbf{z},p^n} \to S^3$ be the Fox completion (cf. e.g. [20]) of the p^n-fold subcover $X_{\mathbf{z},p^n} \to X$ of $X_{\mathbf{z}} \to X$. Then $M_{\mathbf{z},p^n} \to S^3$ is a cyclic branched cover of L associated to $X_{\mathbf{z},p^n} \to X$.

For a fixed prime number p, we put $\widehat{\Lambda}_1 = \mathbb{Z}_p[[T]]$ the ring of formal power series. We can regard Λ_1 as a subring of $\widehat{\Lambda}_1$ by identifying $t = 1 + T$. If $\Delta_{L,\mathbf{z}}(t) \neq 0$, $\Delta_{L,\mathbf{z}}(1+T) \in \widehat{\Lambda}_1$ is uniquely written as

$$\Delta_{L,\mathbf{z}}(1+T) = p^{\mu_{L,\mathbf{z}}} P_{L,\mathbf{z}}(T) U(T)$$

by the p-adic Weierstrass preparation theorem ([27] Theorem 7.3), where $0 \leq \mu_{L,\mathbf{z}} \in \mathbb{Z}$, $U(T) \in \widehat{\Lambda}_1^{\times}$ and $P_{L,\mathbf{z}}(T)$ is a distinguished polynomial of degree $\lambda_{L,\mathbf{z}} = \deg P_{L,\mathbf{z}}(T) \leq \deg' \Delta_{L,\mathbf{z}}(t)$, i.e., a monic polynomial such that $P_{L,\mathbf{z}}(T) \equiv T^{\lambda_{L,\mathbf{z}}} \pmod{p}$. Further, if $\Delta_{L,\mathbf{z}}(t) \in \mathbb{Z}[t]$, then $U(T) \in \mathbb{Z}_p[T]$ ([27] Lemma 7.5).

As pointed out by Mazur [17], the objects analogous to these invariants are as follows. The $\mathbb{Z}_p$-extension k_∞/k is regarded as a tower of cyclic subextensions k_n/k of degree p^n. The Galois group Y_{k_∞} of the maximal unramified abelian pro-p-extension over k_∞ is a module over $\widehat{\Lambda}_1 \simeq \mathbb{Z}_p[[\operatorname{Gal}(k_\infty/k)]] : 1 + T \leftrightarrow \gamma$, where $\operatorname{Gal}(k_\infty/k) = \gamma^{\mathbb{Z}_p}$. The Iwasawa polynomial is defined as $p^{\mu_{k_\infty}} P_{k_\infty}(T) = \Delta^{(0)}_{\widehat{\Lambda}_1}(Y_{k_\infty})$ where $P_{k_\infty}(T)$ is a distinguished polynomial of degree λ_{k_∞}. Then there is an integer ν_{k_∞} satisfying Iwasawa's class number formula (cf. e.g. [27]);

$$v_p(|Cl(k_n)|) = \lambda_{k_\infty} n + \mu_{k_\infty} p^n + \nu_{k_\infty}$$

for all sufficiently large n, where $Cl(k)$ denotes the ideal class group of a number field k which is analogous to $H_1(M; \mathbb{Z})$ of a rational homology 3-sphere M.

infinite cyclic cover $X_{\mathbf{z}} \to X$	$\longleftrightarrow$	$\mathbb{Z}_p$-extension k_∞/k
$\cdots \to M_{\mathbf{z},p^n} \to \cdots \to S^3$	$\longleftrightarrow$	$k \subset \cdots \subset k_n \subset \cdots \subset k_\infty$
Alexander polynomial $\Delta^{(0)}_{\Lambda_1}(H_1(X_{\mathbf{z}}; \mathbb{Z}))$	$\longleftrightarrow$	Iwasawa polynomial $\Delta^{(0)}_{\widehat{\Lambda}_1}(Y_{k_\infty})$

As an analogy of Iwasawa's formula, we have obtained the following theorem (cf. [7] Theorem 5.1.7, [10] Theorem 2.1 and its proof).

Theorem 2.1. (1) *Put* $v = \max\{v_p(z_i) \mid 1 \le i \le r\}$*. For all* $n \ge v$*, we have*

$$|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| = |H_1(M_{\mathbf{z},p^v};\mathbb{Z})| \left| \prod_{\substack{\zeta^{p^n}=1 \\ \zeta^{p^v}\neq 1}} \varDelta_{L,\mathbf{z}}(\zeta) \right|.$$

(2) *If* $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ *for any* n*, there is an integer* $\nu_{L,\mathbf{z}}$ *such that*

$$v_p(|H_1(M_{\mathbf{z},p^n};\mathbb{Z})|) = \lambda_{L,\mathbf{z}} n + \mu_{L,\mathbf{z}} p^n + \nu_{L,\mathbf{z}}$$

for all sufficiently large n*.*

REMARK. The formula of (1) is a direct consequence of Mayberry-Murasugi's formula (cf. [16] [24]). The formula of (2) is induced from (1) by similar arguments to the proof of Theorem 5.1.7 of [7].

By the basic properties of Alexander polynomials, we obtain the following theorem, which decides the parity of $\lambda_{L,\mathbf{z}}$ and the existence of L with prescribed $\lambda_{L,\mathbf{1}}$ and $\mu_{L,\mathbf{1}}$.

Theorem 2.2. (i) *If* $r = 1$*, then* $\lambda_{L,\mathbf{z}} = \mu_{L,\mathbf{z}} = 0$ *and* $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ *for any* n*.*

(ii) *If* $r \ge 2$*, then* $\lambda_{L,\mathbf{z}} \ge r-1$ *and*

$$\lambda_{L,\mathbf{z}} \equiv \begin{cases} r-1 \pmod 2 & \text{if } p \neq 2 \\ \deg' \varDelta_{L,\mathbf{z}}(t) \equiv 1 + \sum_{i=1}^{r} z_i (1 - \sum_{j\neq i} \ell_{ij}) \pmod 2 & \text{if } p = 2 \end{cases}$$

for any $\mathbf{z}$ *such that* $\varDelta_{L,\mathbf{z}}(t) \neq 0$*.*

(iii) *For each* $r \ge 2$ *and for any* $0 \le \ell, m \in \mathbb{Z}$*, there is an* r*-component link* L *such that*

$$\lambda_{L,\mathbf{1}} = r - 1 + 2\ell, \quad \mu_{L,\mathbf{1}} = m$$

and $|H_1(M_{\mathbf{1},p^n};\mathbb{Z})| \neq 0$ *for any* n*.*

Proof. (i) Note that $\mathbf{z} = \pm 1$. Since $\varDelta_{L,\mathbf{z}}(1) = \pm 1$ and $\varPhi_{p^n}(1) = p$, $\varPhi_{p^n}(t)$ does not divide $\varDelta_{L,\mathbf{z}}(t)$ and $\varDelta_{L,\mathbf{z}}(1+T) \in \widehat{\varLambda}_1^{\times}$. Therefore $\lambda_{L,\mathbf{z}} = \mu_{L,\mathbf{z}} = 0$ and $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for any n by Theorem 2.1 (1).

(ii) By Corollary 4.13.2 of [6], $\varDelta_{L,\mathbf{z}}(t)$ is divisible by $(t-1)^{r-1}$, i.e., $P_{L,\mathbf{z}}(T)$ is divisible by T^{r-1}. Therefore $\lambda_{L,\mathbf{z}} \ge r-1$. We may assume that $\varDelta_{L,\mathbf{z}}(t) \in \mathbb{Z}[t]$ and $\varDelta_{L,\mathbf{z}}(0) \neq 0$. Put $\lambda = \lambda_{L,\mathbf{z}}$, $\mu = \mu_{L,\mathbf{z}}$, $P(T) = P_{L,\mathbf{z}}(T)$ and $\dot{T} = (1+T)^{-1} - 1$. By the Torres conditions (cf. [6] etc.), we have $\varDelta_{L,\mathbf{z}}(t) = (-1)^{r-1} t^d \varDelta_{L,\mathbf{z}}(t^{-1})$ where $d = \deg \varDelta_{L,\mathbf{z}}(t) \equiv 1 + \sum_{i=1}^{r} z_i (1 - \sum_{j\neq i} \ell_{ij}) \pmod 2$. Then we have

$$\varDelta_{L,\mathbf{z}}(1+T) = p^{\mu} P(T) U(T) = p^{\mu} (-1)^{r-1} (1+T)^d P(\dot{T}) U(\dot{T})$$

with a distinguished polynomial $P(T) = T^\lambda + \sum_{i=0}^{\lambda-1} c_i T^i \equiv T^\lambda \pmod{p}$ and $U(T) \in \widehat{\varLambda}_1^\times \cap \mathbb{Z}_p[T]$. Since

$$\begin{aligned}(1+T)^\lambda P(\dot{T}) &= (-T)^\lambda + \sum_{i=0}^{\lambda-1} c_i(-T)^i(1+T)^{\lambda-i} \\ &= P(-1)T^\lambda + \sum_{j=0}^{\lambda-1}\sum_{i=0}^{j} c_i(-1)^i\binom{\lambda-i}{j-i}T^j\end{aligned}$$

and the distinguished polynomial $P(T)$ dividing $\varDelta_{L,\mathbf{z}}(1+T)$ in $\widehat{\varLambda}_1$ is unique, then $P(T) = P(-1)^{-1}(1+T)^\lambda P(\dot{T})$ and hence $U(T) = (-1)^{r-1}P(-1)(1+T)^{d-\lambda}U(\dot{T})$. Since $U(0) = (-1)^{r-1}P(-1)U(0)$, we have $(-1)^{r-1} = P(-1) \equiv (-1)^\lambda \pmod{p}$ and $U(T) = (1+T)^{d-\lambda}U(\dot{T})$. Therefore $\lambda \equiv r-1 \pmod{2}$ if $p \neq 2$. Suppose that $p = 2$, and put $u(t) = U(t-1) \in \mathbb{Z}_2[t]$. Since $u(t) = t^{d-\lambda}u(t^{-1})$ and $u(1) = U(0) \not\equiv 0 \pmod{2}$, then $\deg u(t) = d - \lambda$ must be even. Therefore $\lambda \equiv d \pmod{2}$ if $p = 2$.

(iii) Put $\nabla(t) = p^m t^{-\ell}(t-1)^{2\ell} \in \varLambda_1$. Since $\nabla(t) = \nabla(t^{-1})$ and $\deg' \nabla(t)$ is even, there is an r-component link L such that $\varDelta_{L,\mathbf{1}}(t) = (t-1)^{r-1}\nabla(t)$ by [8]. Then $\lambda_{L,\mathbf{1}} = r - 1 + 2\ell$ and $\mu_{L,\mathbf{1}} = m$. By Theorem 2.1 (1), we have $|H_1(M_{\mathbf{1},p^n};\mathbb{Z})| \neq 0$ for all n. $\square$

Remark. We note that $\lambda_{L,\mathbf{1}} \equiv r-1 \pmod{2}$ even if $p = 2$. If p splits completely in k, we have an analogy:

$$T^{r-1} \big| P_{L,\mathbf{1}}(T),\ \lambda_{L,\mathbf{1}} \geq r-1 \quad \longleftrightarrow \quad T^{r_2} \big| P_{k_\infty^{\mathrm{cyc}}}(T),\ \lambda_{k_\infty^{\mathrm{cyc}}} \geq r_2$$

where $k_\infty^{\mathrm{cyc}}/k$ is the cyclotomic $\mathbb{Z}_p$-extension (cf. e.g. [3]). Refer to [1] (resp. [23]) for the existence of k_∞/k with prescribed λ_{k_∞} (resp. μ_{k_∞}).

3. 2-component links

In this section, we consider the Iwasawa invariants of 2-component links.

Lemma 3.1. *For a 2-component link $L = K_1 \cup K_2$ and any $\mathbf{z} = (z_1, z_2)$ with $\gcd(z_1, z_2) = 1$, we have $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for all $0 \leq n \leq v = \max\{v_p(z_1), v_p(z_2)\} = v_p(z_1z_2)$.*

Proof. We may assume that $v = v_p(z_2) \geq v_p(z_1) = 0$. Since the covering space $X_{\mathbf{z},p^n}$ for $n \leq v$ corresponds to the kernel of

$$z_1^{-1} \circ \mathrm{mod} p^n \circ \sigma_{\mathbf{z}} : H_1(X;\mathbb{Z}) \to \mathbb{Z}/p^n\mathbb{Z} : t_1 \mapsto 1 \bmod p^n, t_2 \mapsto 0 \bmod p^n,$$

then $M_{\mathbf{z},p^n}$ is the cyclic branched covering space along K_1. Since $\varDelta_{K_1}(1) = \pm 1 \not\equiv 0 \pmod{p}$, then $\varPhi_{p^n}(t)$ does not divide $\varDelta_{K_1}(t)$ for all $n \geq 1$, and hence $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| = \pm\prod_{\zeta^{p^n}=1}\varDelta_{K_1}(\zeta) \neq 0$ for any $n \leq v$. $\square$

The following theorem shows the relationship between the linking numbers and Iwasawa invariants.

Theorem 3.2. *For a 2-component link L and any $\mathbf{z}$, $\lambda_{L,\mathbf{z}} = 1$, $\mu_{L,\mathbf{z}} = 0$ if and only if $v_p(\ell_{12}) = 0$. Moreover, then $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for any n.*

Proof. By the Torres conditions [26], we have

$$\varDelta_L(t_1,t_2) = \frac{(t_1t_2)^{\ell_{12}}-1}{t_1t_2-1}\varDelta_{K_1}(t_1)\varDelta_{K_2}(t_2) + (t_1-1)(t_2-1)f(t_1,t_2)$$

with some $f(t_1,t_2) \in \varLambda_2$. Then $\varDelta_{L,\mathbf{z}}(1+T)/T \equiv \varDelta_L(1,1) = \pm\ell_{12} \pmod{T}$. Suppose $v_p(\ell_{12}) = 0$. Then $\varPhi_{p^n}(t)$ does not divide $\varDelta_{L,\mathbf{z}}(t)/(t-1)$ for any $n \geq 1$ and $\varDelta_{L,\mathbf{z}}(1+T)/T \in \widehat{\varLambda}_1^{\times}$. By Lemma 3.1 and Theorem 2.1 (1), we have $\lambda_{L,\mathbf{z}} = 1$, $\mu_{L,\mathbf{z}} = 0$ and $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for any n. If $v_p(\ell_{12}) > 0$, then $\varDelta_{L,\mathbf{z}}(1+T)/T \notin \widehat{\varLambda}_1^{\times}$ and hence $\lambda_{L,\mathbf{z}} \geq 2$ or $\mu_{L,\mathbf{z}} \geq 1$. ☐

Since the linking numbers are analogous to the power residue symbols, Theorem 3.2 can be regarded as an analogy of Gold-Sands's theorem (cf. [2], [25] Proposition 2.1). Let k be an imaginary quadratic field in which p splits completely. Then $S = \{\wp_1, \wp_2\}$ and $r_2 = 1$. There is some $\pi_2 \in k$ such that $\wp_2^{h_k} = (\pi_2)$ as a principal ideal. Gold-Sands's theorem implies that $\lambda_{k_\infty} = 1$ and $\mu_{k_\infty} = 0$ (i.e., $Y_{k_\infty} = \mathrm{Gal}(\widetilde{k}/k_\infty) \simeq \mathbb{Z}_p$) for any $\mathbb{Z}_p$-extensions k_∞/k in which both $\wp_1$ and $\wp_2$ ramify if and only if $h_k \not\equiv 0 \pmod{p}$ and $\pi_2^{p-1} \not\equiv 1 \pmod{\wp_1^2}$, i.e., π_2 is not pth power residue modulo $\wp_1^2$.

$$\begin{array}{ccc}
L = K_1 \cup K_2,\ G_L/G_L' \simeq \mathbb{Z}^2 & \longleftrightarrow & S = \{\wp_1,\wp_2\},\ G_S/G_S' \simeq \mathbb{Z}_p^2 \\
|H_1(S^3;\mathbb{Z})| = 1,\ v_p(\ell_{12}) = 0 & \longleftrightarrow & h_k \not\equiv 0 \pmod{p},\ \pi_2^{p-1} \not\equiv 1 \pmod{\wp_1^2} \\
\Updownarrow & & \Updownarrow \\
\lambda_{L,\mathbf{z}} = 1,\ \mu_{L,\mathbf{z}} = 0 & \longleftrightarrow & \lambda_{k_\infty} = 1,\ \mu_{k_\infty} = 0
\end{array}$$

As a generalization of this result, Ozaki [22] proved that if Greenberg's conjecture (cf. §4) holds for $\widetilde{k}/k$ then $\lambda_{k_\infty} = 1$ and $\mu_{k_\infty} = 0$ for all but finitely many k_∞/k. On the other hand, the following theorem implies that if $v_p(\ell_{12}) > 0$, in particular $\ell_{12} = 0$, then $\lambda_{L,\mathbf{z}}$ is not necessary bounded as $\mathbf{z}$ is varied.

Theorem 3.3. (1) *Let $L = K_1 \cup K_2$ be a 2-component link as in Figure 1, where $1 \leq m \in \mathbb{Z}$. Then*

$$\lambda_{L,\mathbf{z}} = 1 + p^{v_p(z_1)} + 3p^{v_p(z_2)}, \quad \mu_{L,\mathbf{z}} = v_p(m)$$

and $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for any n.

(2) *Let $L = C(2a, 2b, -2a)$ be a 2-bridge link in a Conway's notation, where $0 \neq a, b \in \mathbb{Z}$. Then*

$$\lambda_{L,\mathbf{z}} = 2 + p^{v_p(z_1z_2)} + (p^{v_p(a)} - 1)p^{v_p(z_1+z_2)}, \quad \mu_{L,\mathbf{z}} = v_p(b).$$

Moreover, $|H_1(M_{\mathbf{z},p^n};\mathbb{Z})| \neq 0$ for any n if and only if $v_p(z_1z_2) \geq v_p(a)$.

Figure 1: $L = K_1 \cup K_2$, $\ell_{12} = 0$



Proof. For any $1 \leq z \in \mathbb{Z}$, we note that

$$\frac{t^z - 1}{t^{p^{v_p(z)}} - 1} = \sum_{i=0}^{z/p^{v_p(z)}-1} (t^{p^{v_p(z)}})^i \equiv z/p^{v_p(z)} \in \mathbb{Z}_p^\times \pmod{t-1}.$$

(1) By direct computations, one can see that $\varDelta_L(t_1, t_2) = m(t_1 - 1)(t_2 - 1)^3$. Then $\varDelta_{L,\mathbf{z}}(t) = m(t-1)(t^{z_1} - 1)(t^{z_2} - 1)^3$ which is not divisible by $\varPhi_{p^n}(t)$ for all $n > v$. Therefore $|H_1(M_{\mathbf{z},p^n}; \mathbb{Z})| \neq 0$ for any n by Theorem 2.1 (1) and Lemma 3.1. Since

$$\varDelta_{L,\mathbf{z}}(1+T) = p^{v_p(m)} T((1+T)^{p^{v_p(z_1)}} - 1)((1+T)^{p^{v_p(z_2)}} - 1)^3 U(T)$$

with some $U(T) \in \widehat{\varLambda}_1^\times$, we obtain the claim (1).

(2) By [11] etc., we have

$$\varDelta_L(t_1, t_2) = b(t_1 - 1)(t_2 - 1)\frac{(t_1 t_2)^a - 1}{t_1 t_2 - 1}.$$

Put $v = v_p(z_1 z_2)$. Then $\varDelta_{L,\mathbf{z}}(t)$ is divisible by $\varPhi_{p^n}(t)$ if and only if $n \leq v$ or $v_p(z_1 + z_2) < n \leq v_p(a(z_1 + z_2))$. Note that $v_p(z_1 + z_2) = 0$ if $v \neq 0$. By Theorem 2.1 (1) and Lemma 3.1, $|H_1(M_{\mathbf{z},p^n}; \mathbb{Z})| = 0$ for some $n > v$ if and only if $v_p(a) > 0$ and $v < v_p(a(z_1 + z_2))$, i.e., $v_p(a) > v$. Since

$$\varDelta_{L,\mathbf{z}}(1+T) = p^{v_p(b)} T^2((1+T)^{p^v} - 1)\frac{(1+T)^{p^{v_p(a(z_1+z_2))}} - 1}{(1+T)^{p^{v_p(z_1+z_2)}} - 1} U(T)$$

with some $U(T) \in \widehat{\varLambda}_1^\times$, we obtain the claim (2). □

In the case where $p = 2$, Theorem 2.2 (iii) and the following theorem show the existence of a pair of 2-component link L and $\mathbf{z}$ with prescribed $\lambda_{L,\mathbf{z}} \geq 1$ and $\mu_{L,\mathbf{z}} \geq 0$.

Theorem 3.4. *Assume that $p = 2$. For any $0 \leq \ell, m \in \mathbb{Z}$, there is a pair of 2-component link L and $\mathbf{z}$ such that*

$$\lambda_{L,\mathbf{z}} = 2 + 2\ell, \quad \mu_{L,\mathbf{z}} = m$$

and $|H_1(M_{\mathbf{z},2^n}; \mathbb{Z})| \neq 0$ for any n.

Proof. First, we consider the case $\ell \geq 1$. Put

$$\varDelta(t_1,t_2) = 2^m(t_1-1)(t_2-1)(t_1t_2+t_1^{-1}t_2^{-1})^{\max\{0,\ell-2\}}$$

and put $\mathbf{z}=(1,4)$ if $\ell \geq 2$, and $\mathbf{z}=(1,2)$ if $\ell=1$. By a consequence of Bailey's theorem (cf. [6] Corollary 7.4.1, [14]), there is a 2-component link L such that $\varDelta_L = \varDelta(t_1,t_2)$ (and $\ell_{12}=0$). Then

$$\varDelta_{L,\mathbf{z}}(t) = \begin{cases} 2^m(t-1)^2(t^4-1)(t^{10}+1)^{\ell-2}t^{-5(\ell-2)} & \text{if } \ell \geq 2 \\ 2^m(t-1)^2(t^2-1) & \text{if } \ell = 1 \end{cases}$$

and $|H_1(M_{\mathbf{z},2^n};\mathbb{Z})| \neq 0$ for any n by Theorem 2.1 (1) and Lemma 3.1. Since

$$\varDelta_{L,\mathbf{z}}(1+T) = \begin{cases} 2^mT^2((1+T)^4-1)((1+T)^2+1)^{\ell-2}U(T) & \text{if } \ell \geq 2 \\ 2^mT^2((1+T)^2-1)U(T) & \text{if } \ell = 1 \end{cases}$$

with some $U(T) \in \widehat{\varLambda}_1^{\times}$, we have $\lambda_{L,\mathbf{z}} = 2+2\ell$ and $\mu_{L,\mathbf{z}} = m$.

If $\ell = m = 0$, put $L = C(4)$ the 2-bridge link, and put $\mathbf{z}=(1,2)$. Then we have $\varDelta_L = t_1t_2+1$ by [11] and hence $\varDelta_{L,\mathbf{z}}(t) = (t-1)(t^3+1)$. By Theorem 2.1 (1) and Lemma 3.1, $|H_1(M_{\mathbf{z},2^n};\mathbb{Z})| \neq 0$ for any n. Since $\varDelta_{L,\mathbf{z}}(1+T) = T((1+T)^3+1) \equiv T^2(1+T+T^2) \not\equiv 0 \pmod 2$, we have $\lambda_{L,\mathbf{z}} = 2$ and $\mu_{L,\mathbf{z}} = 0$.

Suppose that $\ell = 0$ and $m \geq 1$. Put $\nu(t) = (t^{2^m} - t^{-2^m})/(t-t^{-1})$ and $\beta(t) = t^{-2^m}(t-1)(t^{2^{m+1}-1}-1) \in \varLambda_1$. Since $\nu(t^{-1}) = \nu(t)$ and $\beta(t^{-1}) = \beta(t)$, there are monic polynomials $N(x)$, $B(x) \in \mathbb{Z}[x]$ such that $N(t+t^{-1}) = \nu(t)$ and $B(t+t^{-1}) = \beta(t)$. Then $\deg N(x) = 2^m-1$ and $\deg B(x) = 2^m$. Note that the determinant of the Sylvester matrix Q of $N(x)$ and $B(x)$ is the resultant $\mathrm{Res}_{\mathbb{Z}}(N,B)$ of $N(x)$ and $B(x)$. Let ζ be a primitive 2^{m+1}th root of 1. Since the roots of $N(x)$ are $\zeta^i+\zeta^{-i}$ $(i=1,2,\cdots,2^m-1)$, we have

$$|\mathrm{Res}_{\mathbb{Z}}(N,B)|^2 = \left|\prod_{i=1}^{2^m-1} B(\zeta^i+\zeta^{-i})\right|^2 = \left|\prod_{i=1}^{2^{m+1}-1} \beta(\zeta^i)\right| |\beta(-1)|^{-1} = 2^{2m}$$

and hence $\mathrm{Res}_{\mathbb{Z}}(N,B) = \pm 2^m$. Since each row of Q is contained in the kernel of the surjective homomorphism $\mathbb{Z}^{2^{m+1}-1} \to \mathbb{Z}[x]/(N,B) : (a_{2^{m+1}-2},\cdots,a_1,a_0) \mapsto a_{2^{m+1}-2}x^{2^{m+1}-2}+\cdots+a_1x+a_0 \bmod (N,B)$, we have

$$2^m = \pm\mathrm{Res}_{\mathbb{Z}}(N,B) \in E_{\mathbb{Z}}^{(0)}(\mathbb{Z}[x]/(N,B)) \subset \mathrm{Ann}_{\mathbb{Z}}(\mathbb{Z}[x]/(N,B)).$$

Therefore there exist some $F(x)$, $G(x) \in \mathbb{Z}[x]$ such that $2^m = N(x)F(x)+B(x)G(x) = \nu(t)F(t+t^{-1})+\beta(t)G(t+t^{-1})$. Put $f(t_1,t_2) = F(t_1t_2+t_1^{-1}t_2^{-1})$, $g(t_1,t_2) = G(t_1t_2+t_1^{-1}t_2^{-1})$ and put

$$\varDelta(t_1,t_2) = \frac{(t_1t_2)^{2^{m+1}}-1}{t_1t_2-1}f(t_1,t_2) - (t_1-1)(t_2-1)\frac{(t_1t_2)^{2^{m+1}-1}-1}{t_1t_2-1}g(t_1,t_2).$$

Since $\nu(1) = 2^m$ and $\beta(1) = 0$, then $f(1,1) = F(2) = 1$ and hence $F(t+t^{-1})$ is not divisible by $\Phi_{2^n}(t)$ for all n. By a consequence of Bailey's theorem (cf. [6] Theorem 7.4, [14]), there is a 2-component link L such that $\Delta_L = f(t_1,t_2)^z \Delta(t_1,t_2)$ for some $z \geq 0$ (and $\ell_{12} = 2^{m+1}$). For $\mathbf{z} = (2,-1)$, we have

$$\begin{aligned} \Delta_{L,\mathbf{z}}(t) &= (t^2-1)F(t+t^{-1})^z\big(\nu(t)F(t+t^{-1}) + \beta(t)G(t+t^{-1})\big) \\ &= 2^m(t^2-1)F(t+t^{-1})^z \end{aligned}$$

and $|H_1(M_{\mathbf{z},2^n};\mathbb{Z})| \neq 0$ for any n by Theorem 2.1 (1) and Lemma 3.1. Since $F((1+T)+(1+T)^{-1}) \in \widehat{\Lambda}_1^{\times}$, we have $\lambda_{L,\mathbf{z}} = 2$ and $\mu_{L,\mathbf{z}} = m$. $\square$

REMARK. If $\ell_{12} = 0$, then $\Delta_L(t_1,t_2)$ is divisible by $(t_1-1)(t_2-1)$ and therefore $\lambda_{L,\mathbf{z}} \geq 3$. Since $\Delta_L(1,1) = \pm\ell_{12}$, we have $\mu_{L,\mathbf{z}} \leq v_p(\ell_{12})$ if $\ell_{12} \neq 0$. If $p = 2$ and $\lambda_{L,\mathbf{z}}$ is even, then ℓ_{12} is even and $z_1 + z_2$ is odd by Theorem 2.2 (ii).

4. Greenberg type problem

We shall consider a problem analogous to Greenberg's conjecture. Put the ring $\widehat{\Lambda}_r = \mathbb{Z}_p[[T_1, T_2, \cdots, T_r]]$ of r-variable formal power series. Let G be a pro-p group with a surjective homomorphism $\widehat{\psi} : G \to G/G' = \prod_{i=1}^r \gamma_i^{\mathbb{Z}_p} \simeq \mathbb{Z}_p^r$, and G'' be the commutator subgroup of G'. Then G'/G'' is a module over $\widehat{\Lambda}_r \simeq \mathbb{Z}_p[[G/G']] : 1+T_i \leftrightarrow \gamma_i$, and the completed differential module of G is defined as

$$\mathfrak{A}_G = \bigoplus_{g\in G} \widehat{\Lambda}_r dg \Big/ \big\langle d(g_1g_2) - dg_1 - \widehat{\psi}(g_1)dg_2 \, (g_i \in G)\big\rangle_{\widehat{\Lambda}_r}$$

(cf. [20] §9.3). For the pro-p group G_S, $\mathfrak{X}_S = G_S'/G_S''$ is the S-ramified Iwasawa module which is a finitely generated $\widehat{\Lambda}_{r_2+1}$-module, and we have the completed Crowell exact sequence

$$0 \to \mathfrak{X}_S \xrightarrow{\widehat{\theta}_1} \mathfrak{A}_{G_S} \xrightarrow{\widehat{\theta}_2} \widehat{\Lambda}_{r_2+1} \to \mathbb{Z}_p \to 0$$

where $\widehat{\theta}_1(gG_S'') = dg$ and $\widehat{\theta}_2(dg) = \widehat{\psi}(g) - 1$ (cf. [18] §2.2, [20] §9.4, [21]). Let M_S be the $\widehat{\Lambda}_{r_2+1}$-submodule of $\mathfrak{A}_{G_S}$ generated by $d\tau$ where τ are the elements of the inertia subgroups of G_S. Then $M_S \cap \operatorname{Ker}\widehat{\theta}_2$ is generated by $d\tau$ such that $\tau|_{\widetilde{k}} = 1$, and hence $Y_{\widetilde{k}} = \mathfrak{X}_S/\widehat{\theta}_1^{-1}(M_S \cap \operatorname{Ker}\widehat{\theta}_2)$ is the Galois group of the maximal unramified abelian pro-p-extension of $\widetilde{k}$. Greenberg [5] conjectured that the unramified Iwasawa module $Y_{\widetilde{k}}$ is a pseudonull $\widehat{\Lambda}_{r_2+1}$-module.

The analogue of $Y_{\widetilde{k}}$ is defined in a similar way as follows. Let $\psi : G_L \to G_L/G_L'$ be the natural surjective homomorphism, and put the differential module

$$\mathfrak{A}_L = \bigoplus_{g\in G_L} \Lambda_r dg \Big/ \big\langle d(g_1g_2) - dg_1 - \psi(g_1)dg_2 \, (g_i \in G_L)\big\rangle_{\Lambda_r}$$

of G_L. Let $[\widetilde{m}_i] \in A_L$ be the class of a lift $\widetilde{m}_i$ of m_i with endpoints in $\pi^{-1}(*)$, and M_L the Λ_r-submodule of A_L generated by $[\widetilde{m}_i]$ $(1 \leq i \leq r)$. Since $\mathfrak{A}_L \simeq A_L : d[m_i] \leftrightarrow [\widetilde{m}_i]$, M_L is an analogue of M_S, and we have the Crowell exact sequence

$$0 \to B_L \overset{\theta_1}{\to} A_L \overset{\theta_2}{\to} \Lambda_r \to \mathbb{Z} \to 0$$

(cf. [6] p.70, 78). Thus the analogous module $Y_L = B_L/{\theta_1}^{-1}(M_L \cap \operatorname{Ker}\theta_2)$ is defined.

$$\text{“unramified” link module } Y_L \quad \longleftrightarrow \quad \text{unramified Iwasawa module } Y_{\widetilde{k}}$$

REMARK. It is known that $\operatorname{rank}_{\widehat{\Lambda}_{r_2+1}} \mathfrak{X}_S = r_2$ and $\mathfrak{X}_S$ has no nonzero pseudonull $\widehat{\Lambda}_{r_2+1}$-submodule (cf. [4] [21]). As analogous properties, we know that $\operatorname{rank}_{\Lambda_r} B_L \leq r-1$ and that B_L has no nonzero pseudonull Λ_r-submodule if $\operatorname{rank}_{\Lambda_r} B_L = 0$ (cf. [12] Corollary 7.3.13, [6] Theorem 4.12 (2)).

$$\operatorname{rank}_{\Lambda_r} B_L \leq r-1 \quad \longleftrightarrow \quad \operatorname{rank}_{\widehat{\Lambda}_{r_2+1}} \mathfrak{X}_S = r_2$$

Then the following problem arises as an analogue of Greenberg's conjecture.

Problem. *Is Y_L a pseudonull Λ_r-module ?*

For this problem, we obtain the following criteria and examples. We denote by $\Lambda_{r,J}$ the localization of Λ_r by the multiplicative system $J = \{f \in \Lambda_r \mid f(1,1,\cdots,1) = 1\}$. Note that $\Lambda_{r,J}$ is flat over Λ_r (cf. [15] §7).

Theorem 4.1. (i) *If Δ_L has no prime factor f such that $f(1,1,\cdots,1) = \pm 1$, then Y_L is a pseudonull Λ_r-module.*

(ii) *If $r = 1$, then Y_L is not a pseudonull Λ_1-module.*

Proof. (i) By Levine's theorem [13] (cf. [6] Theorem 4.13), we have $(A_L/M_L) \otimes_{\Lambda_r} \Lambda_{r,J} = 0$ and hence $Y_L \otimes_{\Lambda_r} \Lambda_{r,J} = 0$. Then there exists some $f \in \operatorname{Ann}_{\Lambda_r} Y_L$ such that $f \in J$. On the other hand, $\Delta_L \in E^{(0)}_{\Lambda_r}(B_L) \subset \operatorname{Ann}_{\Lambda_r} B_L \subset \operatorname{Ann}_{\Lambda_r} Y_L$. Since Δ_L and f are relatively prime in Λ_r by the assumption, the divisorial hull of $\operatorname{Ann}_{\Lambda_r} Y_L$ is Λ_r, i.e., Y_L is pseudonull.

(ii) By Theorem 4.12 (1) (2) of [6], $A_L \simeq B_L \oplus \Lambda_1$, and B_L is Λ_1-torsion and has no nonzero pseudonull Λ_1-submodule. Since $(A_L/M_L) \otimes_{\Lambda_1} \Lambda_{1,J} = 0$ by Levine's theorem [13] again, $M_L = \Lambda_1[\widetilde{m}_1] \simeq \Lambda_1$ and hence $M_L \cap \operatorname{Ker}\theta_2 = M_L \cap \theta_1(B_L) = 0$. Therefore $Y_L = B_L$ is not pseudonull. □

EXAMPLE. (1) Let L be a 2-component link as in Figure 1. Then $\Delta_L = m(t_1 - 1)(t_2 - 1)^3$ and hence Y_L is a pseudonull Λ_2-module.

(2) Let $L = C(2p^m, 2b, -2p^m)$ be a 2-bridge link where p is a prime number. Then we have $\Delta_L = b(t_1 - 1)(t_2 - 1)\prod_{n=1}^{m} \Phi_{p^n}(t_1 t_2)$ as in the proof of Theorem

3.3 (2). Note that $\Phi_{p^n}(t_1t_2) \notin J$ since $\Phi_{p^n}(1) = p$. Suppose $\Phi_{p^n}(t_1t_2)$ has a factor $f \in J \cap \mathbb{Z}[t_1, t_2]$ such that $f(0,0) \neq 0$, and put $g = \Phi_{p^n}(t_1t_2)/f$. We denote by $\deg_i$ the degree as a polynomial of one variable t_i. Since $\Phi_{p^n}(t_1)$ is irreducible, we have $g(t_1, 1) = \Phi_{p^n}(t_1)$ and $f(t_1, 1) = 1$. Then $\deg_1 g(t_1, t_2) \geq \deg_1 g(t_1, 1) = \deg_1 \Phi_{p^n}(t_1t_2)$ and hence $\deg_1 f(t_1, t_2) = 0$. Similarly, $\deg_2 f(t_1, t_2) = 0$. This implies that $f = 1$. Therefore Y_L is a pseudonull Λ_2-module.

5. Pro-p link module

We can also regard the pro-p completion $\widehat{G}_L$ of G_L as an analogue of G_S. Then $\widehat{G}_L/\widehat{G}'_L \simeq \mathbb{Z}_p^r$, where $\widehat{G}'_L$ is the commutator subgroup of $\widehat{G}_L$. By identifying $t_i = T_i + 1$, Λ_r and $\Lambda_{r,J}$ are regarded as subrings of $\widehat{\Lambda}_r \simeq \mathbb{Z}_p[[\widehat{G}_L/\widehat{G}'_L]]$. Since $\widehat{\Lambda}_r \simeq \varprojlim \Lambda_{r,J}/\mathfrak{m}^n \simeq \varprojlim \Lambda_r/\mathfrak{m}^n$ where $\mathfrak{m} = (p, T_1, T_2, \cdots, T_r)$ is the maximal ideal, $\widehat{\Lambda}_r$ is flat over Λ_r and over $\Lambda_{r,J}$ (cf. [15] Theorem 8.8). By the following proposition, $\widehat{B}_L = B_L \otimes_{\Lambda_r} \widehat{\Lambda}_r$ is also an analogue of $\mathfrak{X}_S$. Let $\widehat{G}''_L$ be the commutator subgroup of $\widehat{G}'_L$, and put the completed Alexander module $\widehat{A}_L = A_L \otimes_{\Lambda_r} \widehat{\Lambda}_r$.

Proposition 5.1. $\widehat{B}_L \simeq \widehat{G}'_L/\widehat{G}''_L$.

Proof. There is a natural homomorphism $\varphi : \widehat{B}_L \simeq (G'_L/G''_L) \otimes_{\Lambda_r} \widehat{\Lambda}_r \to \widehat{G}'_L/\widehat{G}''_L$. For the natural homomorphism $\widehat{\psi} : \widehat{G}_L \to \widehat{G}_L/\widehat{G}'_L$, the completed differential module $\mathfrak{A}_{\widehat{G}_L}$ of $\widehat{G}_L$ is defined. By [7] (3.1.2) (3.1.6) and [20] Theorem 9.14, we have $\mathfrak{A}_{\widehat{G}_L} \simeq \widehat{A}_L$. Then there is a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \widehat{B}_L \arrow[r, "\theta_1 \otimes id"] \arrow[d, "\varphi"] & \widehat{A}_L \arrow[r, "\theta_2 \otimes id"] \arrow[d, phantom, sloped, "\simeq"] & \Lambda_r \otimes_{\Lambda_r} \widehat{\Lambda}_r \arrow[r] \arrow[d, phantom, sloped, "\simeq"] & \mathbb{Z} \otimes_{\Lambda_r} \widehat{\Lambda}_r \arrow[r] \arrow[d, phantom, sloped, "\simeq"] & 0 \\
0 \arrow[r] & \widehat{G}'_L/\widehat{G}''_L \arrow[r] & \mathfrak{A}_{\widehat{G}_L} \arrow[r] & \widehat{\Lambda}_r \arrow[r] & \mathbb{Z}_p \arrow[r] & 0
\end{tikzcd}$$

with exact rows, where the bottom row is the completed Crowell sequence for $\widehat{G}_L$. Therefore φ is an isomorphism. □

The following proposition is a background of Theorem 3.2.

Proposition 5.2. *If* $r = 2$, *then* $\widehat{B}_L \simeq \widehat{\Lambda}_2/(\Delta_L(1 + T_1, 1 + T_2))$. *Moreover, if* $v_p(\ell_{12}) = 0$, *then* $\widehat{B}_L = 0$.

Proof. Since $E^{(0)}_{\Lambda_2}(B_L) = \Lambda_2\Delta_L$ (cf. [6] Theorem 4.6 (4)) and the presentation matrix of B_L is the same to that of $\widehat{B}_L$, we have $E^{(0)}_{\widehat{\Lambda}_2}(\widehat{B}_L) = \widehat{\Lambda}_2\Delta_L(1 + T_1, 1 + T_2)$. By Theorem 4.7 of [6], $\widehat{B}_L \simeq (B_L \otimes_{\Lambda_2} \Lambda_{2,J}) \otimes_{\Lambda_{2,J}} \widehat{\Lambda}_2$ is a cyclic $\widehat{\Lambda}_2$-module. Therefore $\widehat{B}_L \simeq \widehat{\Lambda}_2/(\Delta_L(1 + T_1, 1 + T_2))$ (cf. [27] p.299 Example (2)). Since $\Delta_L(1, 1) = \pm\ell_{12}$ by the Torres conditions [26], we have $0 \neq \Delta_L(1 + T_1, 1 + T_2) \in \widehat{\Lambda}_2^{\times} \cap \mathrm{Ann}_{\widehat{\Lambda}_2}\widehat{B}_L$ if $v_p(\ell_{12}) = 0$. Then $\widehat{B}_L = 0$. □

Remark. $Y_{\widetilde{k}} = 0$ for the analogous situation (cf. §3). This proposition is also a consequence of Theorem 1.1.6 of [7].

A pro-p version of Greenberg type problem has trivial answer as follows.

Proposition 5.3. $Y_L \otimes_{\Lambda_r} \widehat{\Lambda}_r = 0$.

Proof. By Levine's theorem [13], $Y_L \otimes_{\Lambda_r} \widehat{\Lambda}_r \simeq (Y_L \otimes_{\Lambda_r} \Lambda_{r,J}) \otimes_{\Lambda_{r,J}} \widehat{\Lambda}_r \simeq 0 \otimes_{\Lambda_{r,J}} \widehat{\Lambda}_r \simeq 0$. ☐

This proposition implies that the Iwasawa invariants $\lambda_{L,\mathbf{z}}$, $\mu_{L,\mathbf{z}}$ and $\nu_{L,\mathbf{z}}$ are related to only the structure of the module $M_L \otimes_{\Lambda_r} \widehat{\Lambda}_r$ genetared by meridianal elements.

Acknowledgement. The first author was partially supported by a grant (No. 10801021/a010402) of NSFC, and he was financially supported by Professor Akio Kawauchi during his stay in Japan. The second author was partially supported by Grant-in-Aid for Young Scientists (B) (No. 22740010).

References

[1] S. Fujii, Y. Ohgi and M. Ozaki, *Construction of $\mathbb{Z}_p$-extensions with prescribed Iwasawa λ-invariants*, J. Number Theory **118** (2006), no. 2, 200–207.

[2] R. Gold, *The nontriviality of certain Z_l-extensions*, J. Number Theory **6** (1974), no. 5, 369–373.

[3] R. Greenberg, *On the Iwasawa invariants of totally real number fields*, Amer. J. Math. **98** (1976), no. 1, 263–284.

[4] R. Greenberg, *On the structure of certain Galois groups*, Invent. Math. **47** (1978), no. 1, 85–99.

[5] R. Greenberg, *Iwasawa theory—past and present*, Class field theory—its centenary and prospect (Tokyo, 1998), 335–385, Adv. Stud. Pure Math. **30**, Math. Soc. Japan, Tokyo, 2001.

[6] J. Hillman, *Algebraic invariants of links*, Series on Knots and Everything **32**, World Scientific Publishing Co., Inc., River Edge, NJ, 2002.

[7] J. Hillman, D. Matei and M. Morishita, *Pro-p link groups and p-homology groups*, Primes and knots, 121–136, Contemp. Math. **416**, Amer. Math. Soc., Providence, RI, 2006.

[8] F. Hosokawa, *On ∇-polynomials of links*, Osaka Math. J. **10** (1958), 273–282.

[9] K. Iwasawa, *On Γ-extensions of algebraic number fields*, Bull. Amer. Math. Soc. **65** (1959), 183–226.

[10] T. Kadokami and Y. Mizusawa, *Iwasawa type formula for covers of a link in a rational homology sphere*, J. Knot Theory Ramifications **17** (2008), no. 10, 1199–1221.

[11] T. Kanenobu, *Alexander polynomials of two-bridge links*, J. Austral. Math. Soc. Ser. A **36** (1984), 59–68.

[12] A. Kawauchi, *A survey of knot theory*, Birkhäuser Verlag, Basel, 1996.

[13] J. P. Levine, *Localization of link modules*, Low-dimensional topology (San Francisco, Calif., 1981), 213–229, Contemp. Math. **20**, Amer. Math. Soc., Providence, RI, 1983.

[14] J. P. Levine, *Symmetric presentation of link modules*, Topology Appl. **30** (1988), no. 2, 183–198.

[15] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, Cambridge, 1989.

[16] J. P. Mayberry and K. Murasugi, *Torsion-groups of abelian coverings of links*, Trans. Amer. Math. Soc. **271** (1982), no. 1, 143–173.

[17] B. Mazur, *Remarks on the Alexander polynomial*, unpublished paper, 1963/1964. `http://www.math.harvard.edu/~mazur/papers/alexander_polynomial.pdf`

[18] M. Morishita, *On certain analogies between knots and primes*, J. Reine Angew. Math. **550** (2002), 141–167.

[19] M. Morishita, *Analogies between knots and primes, 3-manifolds and number rings*, Sugaku Expositions **23** (2010), no. 1, 1–30.

[20] M. Morishita, *Knots and Primes - An Introduction to Arithmetic Topology*, Springer, 2012.

[21] T. Nguyen Quang Do, *Formations de classes et modules d'Iwasawa*, Number theory, Noordwijkerhout 1983, 167–185, Lecture Notes in Math. **1068**, Springer, Berlin, 1984.

[22] M. Ozaki, *Iwasawa invariants of $\mathbb{Z}_p$-extensions over an imaginary quadratic field*, Class field theory—its centenary and prospect (Tokyo, 1998), 387–399, Adv. Stud. Pure Math. **30**, Math. Soc. Japan, Tokyo, 2001.

[23] M. Ozaki, *Construction of $\mathbf{Z}_p$-extensions with prescribed Iwasawa modules*, J. Math. Soc. Japan **56** (2004), no. 3, 787–801.

[24] J. Porti, *Mayberry-Murasugi's formula for links in homology 3-spheres*, Proc. Amer. Math. Soc. **132** (2004), no. 11, 3423–3431.

[25] J. W. Sands, *On the nontriviality of the basic Iwasawa λ-invariant for an infinitude of imaginary quadratic fields*, Acta Arith. **65** (1993), no. 3, 243–248.

[26] G. Torres, *On the Alexander polynomial*, Ann. of Math. (2) **57** (1953), 57–89.

[27] L. C. Washington, *Introduction to cyclotomic fields (Second edition)*, Graduate Texts in Mathematics **83**, Springer-Verlag, New York, 1997.

TERUHISA KADOKAMI: Department of Mathematics, East China Normal University, Dongchuan-lu 500, Shanghai 200241, China. `mshj@math.ecnu.edu.cn`

YASUSHI MIZUSAWA: Department of Mathematics, Nagoya Institute of Technology, Gokiso, Showa, Nagoya 466-8555, Japan. `mizusawa.yasushi@nitech.ac.jp`